

Gliadin (Deaminated) Antibody (IgG) Test

DESCRIPTION

Gliadin is one of the major proteins of gluten; the detection of antibodies to gliadin is useful in diagnosing celiac disease. Celiac disease results from an immune-mediated inflammatory process that occurs when consuming foods that have wheat, rye, or barley proteins. The detection of the gliadin antibodies is useful in aiding in the diagnosis or monitoring for certain gluten-sensitive enteropathies, such as celiac disease and dermatitis herpetiformis.

WHY DO I NEED A GLIADIN (DEAMINATED) ANTIBODY (IgG) TEST?

The reasons for you to get this test might be to aid in the diagnosis or monitoring for certain gluten-sensitivities such as celiac disease and dermatitis herpetiformis.

AM I REQUIRED TO FAST?

No, fasting is not required. There are no dietary or medicinal restrictions to take this test.

WHAT IS THE SAMPLE REQUIRED?

A simple venous blood draw

IS A DOCTOR'S ORDER REQUIRED?

No. You do not need to provide a doctor's order to get lab testing done at Any Lab Test Now.

WHEN WILL I GET MY TEST RESULTS?

Test results generally take between 24 to 72 business hours after your specimen is collected.

DO I NEED TO MAKE AN APPOINTMENT?

No. You can walk in for same day testing. We make it easy for you by offering work-friendly hours.

IF I HAVE QUESTIONS ABOUT MY TESTS RESULTS WHO SHOULD I CONTACT?

We recommend you share your results with your primary care physician or with your specialist.

OTHER RELEVANT LAB TESTS

- Celiac Disease Panel
- IgA (Immunoglobulin A)
- Tissue Transglutaminase (tTG)
- Endomysial Antibody